# This Page Is Inserted by IFW Operations and is not a part of the Official Record

### **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

## > IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.






Figure 1

## MODULAR STRATEGY FOR DEVELOPING NON-PEPTIDE PROTEIN KINASE INHIBITORS



Optional side chains.

Figur 2 x-ray structur of (PKA):Mg<sub>2</sub>ATP:pseudosubstrat inhibitor



In Closed Conformation

Figure 3

General Module M<sub>1</sub> Design Features For Binding To The Conserved Protein Kinase Catalytic Region




Figure 4
SUBSTRATE BEHAVIOR FOR BORONIC ACID PKA INHIBITORS 21 & 22





Figure 5

Binding interactions of src substrate
Ac-Ile-Tyr-Gly-Glu-Phe-NH<sub>2</sub> in model src active site.

Figure 6

Design of naphthalene-based src inhibitor scaffold



src:peptide substrate complex

src:naphthalene-based non-peptide inhibitor binding mode

# Figure 7 Design of isoquinoline and indole-based src inhibitor scaffolds

src:isoquinoline-based non-peptide inhibitor binding mode

src:indole-based non-peptide inhibitor binding mode

Figure 8

Figure 9





Figure 12



Figure 13

FIGURE 14







#### FIGURE 15A

#### FIGURE 15B

### Ovarian Tumor N015 Sensitivity To Drugs: Suspended Cell Culture



### Normal Human Fibroblasts Sensitivity To Src Inhibitors: Subconflu nt Cell Culture



KLM 2-31